

Letter to the Editor

The Atrioventricular Canal Defect Is the Congenital Heart Disease Connecting Short Rib-Polydactyly and Oral-Facial-Digital Syndromes

To the Editor:

We read the interesting editorial comment of Neri et al. [1995] pointing out the phenotypic relationship between oral-facial-digital (OFD) and short rib-polydactyly (SRP) syndromes. We agree that a common pathogenetic relation between the two groups does exist. In particular, we think that the study of the anatomic types of congenital heart defects found in both OFD and SRP syndromes may provide further evidence for clinical overlap between the two groups of conditions.

Heart malformations are a cardinal manifestation of the Ellis-van Creveld (EvC) syndrome, the more frequent type being atrioventricular canal defect (AVCD) with common atrium [Lynch et al., 1968; Schinzel, 1983; Marino et al., 1990]. AVC was also reported in the genetic condition signated "single atrium, AVCD-postaxial polydactyly," which is characterized clinically by partial expression of EvC syndrome [Onat, 1994; Digilio et al., 1995a; Levin et al., 1995].

AVCD is the more common heart malformation in patients with Down syndrome [Marino, 1993], and it is frequently found in heterotaxia syndromes [Peoples, 1983]. In addition, our studies have detected a specific association also with monosomy 8p [Marino et al., 1992; Digilio et al., 1993], 3C cranio-cerebello-cardiac syndrome [Digilio et al., 1995b] and Noonan syndrome [Marino et al., 1995]. The specific combination of common atrium and AVCD is rarely diagnosed as isolated malformation in non-syndromic patients. On the contrary, this cardiac defect, frequently with persistent left superior vena cava, is commonly associated with EvC syndrome [Lynch et al., 1968; Schinzel, 1983; Marino et al., 1990] and with heterotaxia syndrome with polysplenia [Peoples, 1983].

Interestingly, the finding of hydrometrocolpos in patients with EvC syndrome [Akoun and Bagard, 1956; Digilio et al., 1996a; Yapar et al., 1996] together

with the report of narrow thoracic cage and short limbs in cases with McKusick-Kaufman (MKK) syndrome [Chitayat et al., 1987] suggests the existence of a pathogenetic link between these autosomal recessive conditions. The MKK syndrome is associated with congenital heart defect in 10% of the cases, AVCD or common atrium being described, to the best of our knowledge, in three of the reported cases [McKusick et al., 1964; McKusick, 1978; Pyeritz et al., 1980].

However, the most interesting evidence is emerging from a review of cardiac malformations in OFD syndromes. Although congenital heart defect is not common in this group, a specific association with AVCD has been detected in patients with OFD syndrome type II [Cordero and Holmes, 1977; Iaccarino et al., 1985; Digilio et al., 1996b]. Furthermore, based on the observation of three patients presenting cardiac defects with oral and digital anomalies characteristic of the OFD syndrome type II but without obvious facial component [Ørstavik et al., 1992; Digilio et al., 1996b], the existence of an "oro-cardio-digital" variant with AVCD has been suggested [Digilio et al., 1996b].

Interestingly, AVCD is also specifically linked with the hydroletharus syndrome, since it was diagnosed in half of the reported patients with congenital heart defect [Salonen et al., 1981; Adetoro et al., 1984; Toriello and Bauserman, 1985].

An attempt at reviewing the known reports of oral-facial-skeletal (OFS) syndromes with AVCD is made in Table I. The nonrandom association of AVCD with EvC, OFD type II, hydroletharus syndromes, and, although less frequently, with other OFS syndromes (Table I) shows that AVCD, with or without common atrium, can represent an additional link between SRP and OFD syndromes.

Clinical variability and genetic heterogeneity of AVCD was demonstrated previously by anatomical and molecular studies [Marino et al., 1990; Wilson et al., 1993; Amati et al., 1995]. Based on the specific association of AVCD with Down and monosomy 8p syndromes, the existence of two "critical regions" for the cardiac defect at chromosome regions 21q22 and 8p23 was suggested [Korenberg et al., 1992; Digilio et al., 1993]. The identification of genes causing oral-facial-skeletal syndromes will likely provide additional candidate loci for genes of AVCD.

*Correspondence to: Maria Cristina Digilio, Servizio di Genetica Medica, Ospedale Bambino Gesù, Piazza S. Onofrio 4, 00165 Rome, Italy.

Received 9 February 1996; Accepted 1 March 1996

TABLE I. Atrioventricular Canal Defect in Oral-Facial-Skeletal Syndromes*

Syndrome	Cardiac defect	References
Ellis-van Creveld	AVCD, AC AVCD, CA	Lynch et al. [1968] Schinzel [1983]; Marino et al. [1990]; Carmi et al. [1992]
Single atrium AVCD/polydactyly	CA AVCD, CA	Onat [1994] Digilio et al. [1995a]; Levin et al. [1995]
Majewski SRP Kaufman-McKusick	AVCD, TA AVCD	Majewski et al. [1971] McKusick et al. [1964]; Peyeritz and O'Neal Humphries [1980] McKusick [1978]
OFD II	CA AVCD, CA, TA AVCD AVCD, AC AVCD, CA	Cordero and Holmes [1977] Iaccarino et al. [1985] Ørstavik et al. [1992] Digilio et al. [1996b]
OFD IIVI	AVCD	Camera et al. [1994]
OFD IV	CA, VSD	Temtamy and McKusick [1978]
OFD VI	AVCD, AC	Gustavson et al. [1971]
Transitional OFD	AVCD	Hingorani et al. [1991]
Pallister-Hall	AVCD	Hall et al. [1980]; Huff and Fernandes [1982]
Hydrolethalus	AVCD	Salonen et al. [1981]; Adetoro et al. [1984]
	AVCD, TrA	Toriello and Bauserman [1985]

* AC, aortic coarctation; AVCD, atrioventricular canal defect; CA, common atrium; OFD, oral-facial-digital; SRP, short rib-polydactyly; TA, truncus arteriosus; TrA, tricuspid atresia; VSD, ventricular septal defect.

REFERENCES

- Adetoro OO, Komolafe F, Anjorin A (1984): Hydrolethalus syndrome in consecutive African siblings. *Pediatr Radiol* 14:422-424.
- Akoun R, Bagard M (1956): Le maladie d'Ellis-van Creveld. *Algérie Méd* 60:769-772.
- Amati F, Mari A, Mingarelli R, Gennarelli M, Digilio MC, Giannotti A, Marino B, Novelli G, Dallapiccola B (1995): Two pedigrees of autosomal dominant atrioventricular canal defect (AVCD): Exclusion from the critical region 8p. *Am J Med Genet* 57:483-488.
- Camera G, Marasini M, Pozzolo S, Camera A (1994): Oral-facial-digital syndrome: Report on a transitional type between the Mohr and Váradí syndromes in a fetus. *Am J Med Genet* 53:196-198.
- Carmi R, Boughman JA, Ferencz C (1992): Endocardial cushion defect: Further studies of "isolated" versus "syndromic" occurrence. *Am J Med Genet* 43:569-575.
- Chitayat D, Hahn SYE, Marion RW, Sachs GS, Goldman D, Hutcheon RG, Weiss R, Cho S, Nitowsky HM (1987): Further delineation of the McKusick-Kaufman hydrometrocolpos-polydactyly syndrome. *Am J Dis Child* 141:1133-1136.
- Cordero JF, Holmes LB (1977): Heart malformation as a feature of the Mohr syndrome. *J Pediatr* 1977;91:683-684.
- Digilio MC, Giannotti A, Marino B, Dallapiccola B (1993): Atrioventricular canal and 8p-syndrome. *Am J Med Genet* 47:437-438.
- Digilio MC, Marino B, Giannotti A, Dallapiccola B (1995a): Single atrium, atrioventricular canal/postaxial hexodactyly indicating Ellis-van Creveld syndrome. *Hum Genet* 96:251-253.
- Digilio MC, Marino B, Giannotti A, Mingarelli R, Dallapiccola B (1995b): Atrioventricular canal and 3C (cranio-cerebello-cardiac) syndrome. *Am J Med Genet* 58:97-98.
- Digilio MC, Marino B, Giannotti A, Dallapiccola B (1996a): Atrioventricular canal defect and postaxial polydactyly indicating phenotypic overlap of Ellis-van Creveld and Kaufman-McKusick syndromes. *Pediatr Cardiol* (in press).
- Digilio MC, Marino B, Giannotti A, Dallapiccola B (1996b): Orocardio-digital syndrome: An oral-facial-digital type II-variant associated with atrioventricular canal. *J Med Genet* 33:416-418.
- Gustavson KH, Kreuger A, Petersson PO (1971): Syndrome characterized by lingual malformation, polydactyly, tachypnea, and psychomotor retardation (Mohr syndrome). *Clin Genet* 2:261-266.
- Hall JG, Pallister PD, Clarren SK, Beckwith JB, Wiglesworth FW, Fraser FC, Cho S, Benke PJ, Reed SD (1980): Congenital hypothalamic hamartoblastoma, hypopituitarism, imperforate anus, and postaxial polydactyly. A new syndrome? Part I: Clinical, causal, and pathogenetic considerations. *Am J Med Genet* 7:47-74.
- Hingorani SR, Pagon RA, Shepard TH, Kapur RP (1991): Twin fetuses with abnormalities that overlap with three midline malformation complexes. *Am J Med Genet* 41:230-235.
- Huff DS, Fernandes M (1982): Two cases of congenital hypothalamic hamartoblastoma, polydactyly, and other congenital anomalies (Pallister-Hall syndrome). *N Engl J Med* 306:430-431.
- Iaccarino M, Lonardo F, Giugliano M, Della Bruna M (1985): Prenatal diagnosis of Mohr syndrome by ultrasonography. *Prenat Diagn* 5:415-418.
- Korenberg JR, Bradley C, Distèche CM (1992): Down syndrome: Molecular mapping of the congenital heart disease and duodenal stenosis. *Am J Hum Genet* 50:294-302.
- Levin SE, Dansky R, Milner S, Benatar A, Govendrageloo K, du Plessis J (1995): Atrioventricular septal defect and type A postaxial polydactyly without other major associated anomalies: A specific association. *Pediatr Cardiol* 16:242-246.
- Lynch JI, Perry LW, Takakuwa T, Scott LP (1968): Congenital heart disease and chondroectodermal dysplasia. Report of two cases, one in a Negro. *Am J Dis Child* 115:80-87.
- Majewski F, Pfeiffer RA, Lenz W, Muller R, Feil G, Seiler R (1971): Polysyndactylie, verkürzte Gliedmassen und Genitalfehlbildungen: Kennzeichen eines selbständigen Syndroms? *Z Kinderheilk* 111:118-138.
- Marino B, Vairo U, Corno A, Nava S, Guccione P, Calabrò R, Marcelletti C (1990): Atrioventricular canal in Down syndrome. Prevalence of associated cardiac malformations compared with patients without Down syndrome. *Am J Dis Child* 144:1120-1122.
- Marino B, Reale A, Giannotti A, Digilio MC, Dallapiccola B (1992): Nonrandom association of atrioventricular canal and del(8p) syndrome. *Am J Med Genet* 42:424-427.
- Marino B (1993): Congenital heart disease in patients with Down's syndrome: Anatomic and genetic aspects. *Biomed Pharmacother* 47:197-200.
- Marino B, Gagliardi MG, Digilio MC, Polletta B, Grazioli S, Agostino D, Giannotti A, Dallapiccola B (1995): Noonan syndrome: Structural

- abnormalities of the mitral valve causing subaortic obstruction. *Eur J Pediatr* 154:949–952.
- McKusick VA, Bauer RL, Koop CE, Scott RB (1964): Hydrometrocolpos as a simply inherited malformation. *JAMA* 189:813–816.
- McKusick VA (1978): The William Allen Memorial Lecture. Genetic nosology. Three approaches. *Am J Hum Genet* 30:105–122.
- Neri G, Gurrieri F, Genuardi M (1995): Oral-facial-skeletal syndromes. *Am J Med Genet* 59:365–368.
- Onat T (1994): Postaxial hexadactyly and single atrium: A new syndrome? *Hum Genet* 94:104–106.
- Östvik KH, Lindenmann R, Solberg LA, Foerster A, Sorland SJ (1992): Congenital heart defects, hamartomas of the tongue and polysyndactyly in a sister and brother. *Clin Genet* 42:19–21.
- Peoples WM, Moller JH, Edwards JE (1983): Polysplenia: A review of 146 cases. *Pediatr Cardiol* 4:129–138.
- Pyeritz RE, O'Neal Humphries J (1980): Partial endocardial cushion defect and persistence of the left superior vena cava draining into the left atrium in a 34-year-old man with features of the Kaufman and Marfan syndromes. *Johns Hopk Med J* 146:28–32.
- Salonen R, Herva R, Norio R (1981): The hydroletharus syndrome: Delineation of a “new,” lethal malformation syndrome based on 28 patients. *Clin Genet* 19:321–330.
- Schinzel AA (1983): Cardiovascular defects associated with chromosomal aberrations and malformation syndromes. *Progr Med Genet* 5:303–379.
- Temtam S, McKusick VA (1978) The genetics of hand malformations, 2nd Ed. New York: Alan R. Liss, Inc., for the National Foundation—March of Dimes. BD-OAS XIV(3):431–437.
- Toriello HV, Bauserman SC (1985): Bilateral pulmonary agenesis: association with the hydroletharus syndrome and review of the literature from a developmental field perspective. *Am J Med Genet* 21:93–103.
- Wilson L, Curtis A, Korenberg JR, Schipper RD, Allan L, Chenevix-Trench G, Stephenson A, Goodship J, Burn J (1993): A large, dominant pedigree of atrioventricular septal defect (AVSD): Exclusion from the Down syndrome critical region on chromosome 21. *Am J Hum Genet* 53:1262–1268.
- Yapar EG, Ekici E, Aydogdu T, Senses E, Gökmen O (1996): Diagnostic problems associated with McKusick Kaufman syndrome (mucometrocolpos, polydactyly and congenital heart disease) with skeletal dysplasia. *Am J Med Genet* 66:343–346.

Maria Cristina Digilio*

Bruno Marino

Aldo Giannotti

Departments of Medical Genetics

and Pediatric Cardiology

Bambino Gesù Hospital

Rome, Italy

Bruno Dallapiccola

Chair of Human Genetics

Public Health and Cell Biology

University of Tor Vergata

Rome, Italy